

Determination of Higgs Couplings at LHC \oplus ILC

Sven Heinemeyer, CERN

SLAC, 03/2005

based on collaboration with

K. Desch, M. Dürrssen, H. Logan, D. Rainwater, G. Weiglein, D. Zeppenfeld

1. Motivation
2. Higgs coupling determination at the LHC
3. Higgs coupling determination at the ILC
4. LHC \oplus ILC
5. Conclusions

1. Motivation

Higgs field in the SM:

$$\Phi = \begin{pmatrix} 0 \\ v + H \end{pmatrix} \quad (\text{unitary gauge})$$


H : elementary scalar field, Higgs boson

Lagrange density: $\mathcal{L}_{\text{Higgs}} = (D_\mu \Phi)^\dagger (D^\mu \Phi) - V(\Phi)$

Gauge invariant coupling to gauge fields

\Rightarrow mass terms for gauge bosons and fermions

1.) $VV\Phi\Phi$ coupling:




\Rightarrow VV mass terms

\Rightarrow triple/quartic couplings to gauge bosons


$$g_2^2 v^2 / 2 \equiv M_W^2, \quad (g_1^2 + g_2^2) v^2 / 2 \equiv M_Z^2 \quad \Rightarrow \text{coupling} \propto \text{masses}$$

2.) fermion mass terms: Yukawa couplings



$$m_f = v g_f \Rightarrow \text{coupling} \propto \text{masses}$$

3.) mass of the Higgs boson: self coupling



$$\lambda = M_H^2/v$$

$$M_H = v\sqrt{\lambda} \quad \text{free parameter}$$

→ last unknown parameter of the SM


⇒ establish Higgs mechanism \equiv find the Higgs \oplus measure its couplings

Higgs coupling measurements is one of the main tasks of future colliders

2. Higgs coupling determination at the LHC

[M. Dürrsen, S.H., H. Logan, D. Rainwater, G. Weiglein, D. Zeppenfeld '04]

Higgs production at the LHC:



gluon fusion: $gg \rightarrow H$

weak boson fusion (WBF):
 $q\bar{q} \rightarrow q'\bar{q}'H$

top quark associated
production: $gg, q\bar{q} \rightarrow t\bar{t}H$

weak boson associated
production: $q\bar{q}' \rightarrow WH, ZH$

Some LHC specifics:

No LHC analogue to recoil method at LEP/LC: $e^+e^- \rightarrow ZH, Z \rightarrow e^+e^-, \mu^+\mu^-$

\Rightarrow no total measurement of Higgs production cross section

QCD backgrounds \Rightarrow not all decay modes accessible, e.g. $H \rightarrow b\bar{b}$

Measurement of $\sigma \times \text{BR}$: narrow width approximation:

$$\Rightarrow \sigma(H) \times \text{BR}(H \rightarrow d_1 d_2) = \frac{\sigma(H)^{\text{SM}}}{\Gamma_{\text{prod}}^{\text{SM}}} \times \frac{\Gamma_{\text{prod}} \Gamma_{\text{decay}}}{\Gamma_{\text{tot}}}$$

Observation of different channels (or upper bound from non-observation)

\Rightarrow information on combinations of $\Gamma_b, \Gamma_\tau, \Gamma_W, \Gamma_Z, \Gamma_g, \Gamma_\gamma, Y_t^2$

\Rightarrow Determination of ratios of partial width via global fit

[M. Dührssen '03]

\Rightarrow Additional theoretical assumptions needed for absolute determination of partial widths

Strategy: mild theoretical assumptions

→ consider **general multi-Higgs-doublet model**
(w/o additional Higgs singlets)
(\Rightarrow including e.g. MSSM)

$\Rightarrow HVV$ coupling bounded from above by SM value, $\Gamma_V \leq \Gamma_V^{\text{SM}}$, $V = W, Z$
 \Rightarrow upper bound on Γ_V

Observation of Higgs production

\Rightarrow lower bound on production couplings
lower bound on total width Γ_{tot}

Observation of $H \rightarrow VV$ in WBF

\Rightarrow determines $\Gamma_V^2/\Gamma_{\text{tot}}$
 \Rightarrow determines upper bound on Γ_{tot}

\Rightarrow Absolute determination of Γ_{tot} and Higgs couplings via global fit

\Rightarrow nearly model independent analysis

Estimate of errors:

Statistical errors:

Assume **SM rates** for production and decay in each luminosity scenario

Systematic errors:

- 5% luminosity error
- uncertainties on reconstruction: identification of leptons: 2%
identification of photons: 2%
identification of b quarks: 3%
- forward tagging/veto jets: 5%
- error propagation for background determination from side-band analyses:
from 0.1% ($H \rightarrow \gamma\gamma$) to 5% ($H \rightarrow WW^*, H \rightarrow \tau^+\tau^-$)
- theoretical and parametric uncertainties for Higgs production:
 ggH : 20%, $t\bar{t}H$: 15%, WH, ZH : 7%, WBF: 4%
- theoretical and parametric uncertainties on Higgs decays:
1% (as a future expectation)

⇒ log likelihood function based on statistical and systematic errors

Decay channels considered:

- $H \rightarrow W^{+(*)}W^{-(*)} \rightarrow l^+l^- + p_{\text{T,miss}}$
- $H \rightarrow Z^{(*)}Z^{(*)}$
- $H \rightarrow \gamma\gamma$
- $H \rightarrow \tau^+\tau^-$
- $t\bar{t}H, H \rightarrow b\bar{b}$


Actual analysis assumptions (even less restrictive):

$$g_{HVV}^2 \leq 1.05 \times g_{HVV,SM}^2, \quad V = W, Z$$

5% margin to allow for

- theoretical uncertainties in translation of partial widths to g_{HVV}^2
- small admixtures of exotic states (triplets, ...)
- Allow for additional particles contributing to $H \rightarrow \gamma\gamma$ and $gg \rightarrow H$
(\Rightarrow fitted by pos. /neg. additional partial width to $H \rightarrow \gamma\gamma$ and $gg \rightarrow H$)
- Allow for additional Higgs decay width
(\Rightarrow fitted by additional partial width)

Constraints on extra partial widths:



Detection of SM rates
 \Rightarrow constraints on widths:

$2 * 300 + 2 * 100 \text{ fb}^{-1}$ scenario:



$$\Delta\Gamma_\gamma \leq 0.2 \times \Gamma_\gamma^{\text{SM}}$$

$$\Delta\Gamma_g \leq 0.4 \times \Gamma_g^{\text{SM}}$$

$$\Delta\Gamma_{\text{inv}} \leq 0.2 \times \Gamma_{\text{tot}}^{\text{SM}}$$

\Rightarrow restrictions on new physics

Relative precisions for partial and total widths: two scenarios



Observations:

low luminosity scenario: $2 * 30 \text{ fb}^{-1}$:

for a light Higgs: results significantly worse as compared to higher luminosity scenario

high(er) luminosity scenario: $2 * 300 + 2 * 100 \text{ fb}^{-1}$:

- typical accuracies of 20-30% for $m_H \leq 150 \text{ GeV}$
- 10% accuracies for HVV couplings above WW threshold

high luminosity scenario: $2 * 300 \text{ fb}^{-1}$:


significant improvement over $2 * 300 + 2 * 100 \text{ fb}^{-1}$ only in $H\tau\tau$ coupling (WBF crucial for $H \rightarrow \tau^+\tau^-$)

Systematic errors contribute up to half of the total error, especially at high luminosity

3. Higgs coupling determination at the ILC

[TESLA TDR '01] [Abe et al. '01] [Abe et al. '01] [T. Barklow '03]

Higgs production at the ILC:



Higgs-strahlung:



weak boson fusion (WBF):



Some ILC specifics:

recoil method: $e^+e^- \rightarrow ZH, Z \rightarrow e^+e^-, \mu^+\mu^-$

⇒ total measurement of Higgs production cross section

⇒ **NO** additional theoretical assumptions needed for absolute determination of partial widths

⇒ all observable channels can be measured with high accuracy

Some ILC results ($500 \text{ fb}^{-1} @ \sqrt{s} = 350 \text{ GeV}$):

$$\delta M_H \approx 50 \text{ MeV}$$

$$\delta g_{ZZH} \approx 2.5\%, \quad \delta g_{WWH} \approx 2 - 5\%$$

$$\delta g_{Hb\bar{b}} \approx 1 - 2\% \text{ (for } M_H \lesssim 150 \text{ GeV)}$$

However: **No good determination of $g_{Ht\bar{t}}$ and $g_{H\gamma\gamma}$**

$$\delta g_{Ht\bar{t}} = ?, \quad \delta g_{H\gamma\gamma} = 23\% \text{ (for } M_H = 120 \text{ GeV)}$$

4. LHC \oplus ILC

Idea: use data from the 350 GeV ILC
to replace theory assumptions in LHC analysis

\Rightarrow better determination of $g_{Ht\bar{t}}$ and $g_{H\gamma\gamma}$
(in a model independent analysis)

Existing result:

[K. Desch, M. Schumacher '04]

LHC: $t\bar{t}H \rightarrow t\bar{t} WW^*, t\bar{t} b\bar{b}$ (300 fb⁻¹)

ILC: BR($H \rightarrow WW^*$), BR($H \rightarrow b\bar{b}$) (500 fb⁻¹ @ $\sqrt{s} = 500$ GeV)


\Rightarrow fit for $g_{Ht\bar{t}}$

\rightarrow T

\Rightarrow determination of $g_{Ht\bar{t}}$ to 15%

Fit for $g_{Ht\bar{t}}$:

[K. Desch, M. Schumacher '04]



New result:

→ Use more ILC input and all LHC channels in a combined fit

Inputs from the ILC to the fit:

– M_H

– $\sigma_{\text{tot}}(e^+e^- \rightarrow HZ)$



– $\sigma_{\text{tot}}(e^+e^- \rightarrow HZ) \times \text{BR}(H \rightarrow X)$ ($X = b\bar{b}, \tau^+\tau^-, gg, WW^*$)

– $\sigma_{\text{tot}}(e^+e^- \rightarrow \nu\bar{\nu}H) \times \text{BR}(H \rightarrow b\bar{b})$


⇒ (hopefully) better determination of $g_{Ht\bar{t}}, g_{H\gamma\gamma}$
(in a model independent way!)

WARNING: results still preliminary

Compare old and new results for $g_{Ht\bar{t}}$:



New results for $g_{Ht\bar{t}}$:



Note: we show Δg^2



[K. Desch, M. Schumacher '04] show Δg

drastic improvement in all channels
 due to ILC input
 (often ILC precision)


⇒ determination of $g_{Ht\bar{t}}$ to 11-14%

⇒ somewhat better than other analysis

Compare old and new results for $g_{H\gamma\gamma}$:



New results for $g_{H\gamma\gamma}$:



⇒ determination of $g_{H\gamma\gamma}$ to $\sim 7 - 8\%$

⇒ ILC input helps at lower M_H values

5. Conclusions

- Higgs coupling determination at the LHC:

coupling determination necessary to establish the Higgs mechanism

→ nearly model independent analysis

⇒ coupling determination down to 20-40%

- Idea: use ILC input to overcome theory assumptions

- ILC input from 500 fb^{-1} @ $\sqrt{s} = 350 \text{ GeV}$

(→ ILC will determine nearly all couplings model independent with a high accuracy)

- However: no good ILC ($\sqrt{s} \lesssim 500 \text{ GeV}$) precision for $g_{Ht\bar{t}}$ and $g_{H\gamma\gamma}$

- New (preliminary) result:

$\Delta g_{Ht\bar{t}} = 11 - 14\%$ (old LHC/ILC analysis: 13-19%)

$\Delta g_{H\gamma\gamma} = 7 - 8\%$ (old LHC analysis: 8-30%)